

# Comparison of climate time series produced by General Circulation Models and by observed data on a global scale

A.M. Filippidou, A. Andrianopoulos, C. Argyrakis, L. E. Chomata, V. Dagalaki, X. Grigoris, T. S. Kokkoris, M. Nasioka, K. A. Papazoglou, S. M. Papalexiou, H. Tyralis and D. Koutsoyiannis  
Department of Water Resources and Environmental Engineering, National Technical University of Athens (itia.ntua.gr/1440)

## 1. Abstract


Outputs of General Circulation Models (GCMs) for precipitation are compared with time series produced from observations. Comparison is made on global and hemispheric spatial scale and on annual time scale. Various time periods are examined, distinguishing periods before and after publishing of model outputs. Historical climate time series are compared with the outputs of GCMs for the 20th century and those for the A1B, B1 and A2 emission scenarios for the 21st century. Several indices are examined, i.e. the estimated means, variances, Hurst parameters, cross-correlations etc.

Acknowledgement: This research is conducted within the frame of the undergraduate course "Stochastic Methods in Water Resources" of the National Technical University of Athens (NTUA). The School of Civil Engineering of NTUA provided moral support for the participation of the students in the Assembly.

## 2. Introduction


- Precipitation data at the annual time scale are compared to output of GCMs for the 20C3M, A1B, B1, A2 scenarios (Hegerl et al. 2003; IPCC 2000; IPCC 2007; IPCC-TGCI A 1999; Leggett et al. 1992).
- We examine:
  - land and land combined with sea regions;
  - the Southern, the Northern Hemisphere and the globe.
- The precipitation data sets are the GPCP Version 2.2 (Adler et al. 2003), the CRU TS3.10 .01 (Harris et al. 2013) and the GPCC Version 6.0 (Schneider et al. 2011).
- The GCM outputs for the IPCC Fourth Assessment Report (AR4) (IPCC 2007) are used for the comparison.
- Data and their integration on the above regions were obtained from the KNMI Climate Explorer web site (climexp.knmi.nl).
- The Hurst parameter, the standard deviation, the linear trend and the cross-correlation for corresponding time periods are estimated and compared.
- Previous similar studies have been performed for smaller regions e.g. Anagnostopoulos et al. (2010), Koutsoyiannis et al. (2008). It was shown that the GCMs for the 20C3M scenario failed to reproduce the past climate.
- Here almost 10 years after the preparation of the AR4, the predictions of the A1B, B1, A2 scenarios are compared to the obtained data sets in addition to the comparison with the 20C3M scenario.

## 3. 20C3M scenario for the land and sea regions




The Figures represent the annual precipitation rate per year. Notice the difference between the observed data (GPCP data set) and the GCM in terms of their means and standard deviations.

## 4. 20C3M scenario for the land regions




The Figures represent the annual precipitation rate per year. Notice the difference between the observed data (CRU and GPCC data sets) and the GCM in terms of their means and standard deviations. Notice also the difference in the means of CRU and GPCC data sets.

## 5. A1B scenario for the land and sea regions




The Figures represent the annual precipitation rate per year. Notice the difference between the observed data (GPCP data set) and the GCM in terms of their means and standard deviations. Notice also that the GCMs predicted increase of the precipitation, which has not been verified by the data.

## 6. A1B scenario for the land regions




The Figures represent the annual precipitation rate per year. Notice the difference between the observed data (CRU and GPCC data sets) and the GCM in terms of their means and standard deviations. Notice also that the GCMs predicted increase of the precipitation, which has not been verified by the data except in GPCC data for the Northern Hemisphere.

## 7. Hurst parameter comparison




Green lines indicate differences between Hurst parameter estimates up to 0.2. Almost half of the models are out of this range.

## 8. Standard deviation estimate comparison




Green lines indicate differences between standard deviations estimates up to 10 mm/year. Almost 75% of the models are out of this range. Estimated standard deviations of models tend to be smaller.

## 9. Linear trend comparison



Green lines indicate differences between linear trend estimates up to 4 mm/year². The 20C3M scenario succeeded in representing the observed data, in which the trend is virtually 0. However almost 50% of the models for the A1B, B1, A2 scenarios are out of this range.

## 10. Cross-correlation estimate comparison



Cross-correlations for lag-zero estimates histograms between the annual values of the observed data sets and the outputs of GCMs. The cross-correlations for the 20C3M are concentrated around zero, whereas for the other models they are uniformly distributed in the whole domain.

## 11. Conclusions

- The GCMS do not reproduce:
  - the mean,
  - the standard deviation,
  - the Hurst parameter, and
  - the linear trendof the observed precipitation.
- The GCMS do not correlate adequately with the observed precipitation.
- Furthermore, there are discrepancies among the different data sets of global and hemispheric precipitation. In particular the estimated statistical parameters of the CRU and GPCC data sets (observed precipitation) have differences at the annual time scale. However, they correlate adequately as shown in the table below.

	Southern hemisphere		Northern hemisphere		Global	
	CRU	GPCC	CRU	GPCC	CRU	GPCC
Hurst parameter	0.75	0.53	0.76	0.56	0.83	0.61
Standard deviation (mm/year)	47.09	58.39	16.21	23.34	20.21	22.99
Linear trend (mm/year²)	0.39	0.12	0.12	0.22	0.21	0.19
Cross-correlation	0.84		0.67		0.78	

## 12. References

Adler RF, Huffman GJ, Chang A, Ferraro R, Xie PP, Janowiak J, Rudolf B, Schneider U, Curtis S, Bolvin D, Gruber A, Susskind J, Arkin P (2003) The Version 2 Global Precipitation Climatology Project (GPCP) Monthly Precipitation Analysis (1979-Present). Journal of Hydrometeorology 4(6):1147-1167. doi:10.1175/JHM2-7541(2003)04<1147:GPCP2.0.CO;2

Anagnostopoulos GC, Koutsoyiannis D, Christofides A, Elstratidis A, Mamassis N (2010) A comparison of local and aggregated climate model outputs with observed data. Hydrological Sciences Journal 55(7):1094-1110. doi:10.1080/02626667.2010.513518

Harris I, Jones PD, Osborn TJ, Lister DH (2013) Updated high-resolution grids of monthly climatic observations – the CRU TS3.10 Dataset. International Journal of Climatology 34(3):623-642. doi:10.1002/joc.3711

Hegerl GC, Mehl G, Covey C, Latif M, McAvaney B, Stouffer R (2003) 20C3M: CMIP collecting data from 20th century coupled model simulations. CLIVAR Exchanges 26 8(1)

IPCC (2000) Special Report on Emissions Scenarios. Nakicenovic N, Swart RJ (eds). Cambridge University Press, Cambridge, United Kingdom

IPCC (2007) Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (eds). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA

IPCC-TGCI A (1999) Guidelines on the use of scenario data for climate impact and adaptation assessment. Carter TR, Hulme M, Lal M (prepared by). Intergovernmental Panel on Climate Change, Task Group on Scenarios for Climate Impact Assessment

Koutsoyiannis D, Elstratidis A, Mamassis N, Christofides A (2008) On the credibility of climate predictions. Hydrological Sciences Journal 53(4):671-684. doi:10.1623/hysj.53.4.671

Leggett J, Pepper WJ, Swart RJ (1992) Emissions scenarios for the IPCC: an update. In Houghton TJ, Berger JO, Callander BA, Varney SK (eds) Climate Change 1992: The Supplementary Report to the IPCC Scientific Assessment. Cambridge University Press, Cambridge, UK, pp 69-95

Schneider U, Becker A, Finger P, Meyer-Christoffer A, Rudolf B, Ziese M (2011) GPCC Full Data Reanalysis Version 6.0 at 0.5°. Monthly Land-Surface Precipitation from Rain-Gauges built on GTS-based and Historic Data. doi:10.5676/DWD\_GPCC/FD\_M\_V6\_050