


# The role of teleconnections in extreme (high and low) precipitation events: The case of the Mediterranean region

Y. Markonis, D. Koutsoyiannis and S. M. Papalexiou, Department of Water Resources and Environmental Engineering, National Technical University of Athens (itia.ntua.gr/1324)


## 1. Abstract

During the last years large-scale climatic indices, such as North Atlantic Oscillation (NAO) and El-Niño Southern Oscillation (ENSO), have been used to describe a certain portion of climatic variability in different temporal and spatial scales. In this context, the climate in the Mediterranean region has been mainly correlated with the NAO index, while there is also some evidence for seasonal associations with the Dynamic Indian Monsoon Index (DIMI) during the summer, and the Siberian High during the winter. Here, we explore the possible links between extreme (high and low) precipitation events in the Mediterranean basin and several large-scale climatic indices, such as these mentioned above and also East Atlantic Pattern, Scandinavia Pattern, Polar/Eurasia Pattern, Mediterranean Oscillation Index, West Africa Monsoon Index and Siberian High. In order to achieve that, we use precipitation data from the Global Historical Climatology Network (GHCN) and index data from National Oceanic and Atmospheric Administration (NOAA).


## 4. Climatic indices




## 7. Monthly extremes (cont.)




## 10. Discussion of the results




## 2. Rainfall data



## 5. Daily maxima




## 8. Winter and annual extremes




## 11. Moving to larger scales




## 3. Homogeneous regions (with respect to rainfall)



## 6. Monthly extremes



## 9. Winter extremes



## 12. Conclusions

The general atmospheric circulation is linked to both high and low extreme precipitation events in the Mediterranean region. This is demonstrated by the two tables below (red is used for positive index values and blue for negative). Furthermore, the main conclusions drawn by our analysis are:

- The correlation becomes higher, as scale increases. More specifically, daily precipitation maxima demonstrate the least correlation, due to a number of reasons including: the scale difference between the indices and the rain event, the convective storms which are best described by synoptic circulation, and smaller scale cyclogenesis at the Mediterranean Sea (e.g. the Gulf of Genoa).
- Using the annual indices could be misleading, even if they refer to the hydrological year, because of their enhanced inter-annual variability. Hence the annual scale is described better if the indices are aggregated to the winter scale.
- The atmospheric circulation has a more direct effect to precipitation minima, as a consequence of the blocking of wet aeral masses towards the Mediterranean. A question left to be answered is whether it also affects the regional cyclogenesis.

Index	Maximum precipitation						Minimum precipitation					
	Iberian Pen.	Central Med.	Balkan Pen.	Mid. East	NC. Africa	W. Africa	Iberian Pen.	Central Med.	Balkan Pen.	Mid. East	NC. Africa	W. Africa
NAO	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year
EA	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year
SCAND	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year
PEU	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year
MOI	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year	month, year
ENSO	year	year	year	year	year	year	year	year	year	year	year	year
SHI	year	year	year	year	year	year	year	year	year	year	year	year
WAMI	year	year	year	year	year	year	year	year	year	year	year	year
DIMI	year	year	year	year	year	year	year	year	year	year	year	year

References:

Haylock, M. R., & Goodess, C. M. (2004). Interannual variability of European extreme winter rainfall and links with mean large-scale circulation. *International Journal of Climatology*, 24(6), 759-776. doi:10.1002/joc.1033

Hoerling, M., Eischeid, J., Pielke, J., Quan, X., Zhang, T., & Pegen, P. (2012). On the Increased Frequency of Mediterranean Drought. *Journal of Climate*, 25(6), 2146-2161. doi:10.1175/JCLI-D-11-00296.1

Hurrell, J. W. (1995). Decadal trends in the North Atlantic oscillation: Regional temperatures and precipitation. *Science*, 269, 676-679

Trigo, R. M., Osborn, T. J., & Corte-Real, J. M. (2002). The North Atlantic Oscillation influence on Europe: climate impacts and associated physical mechanisms. *Climatic Research*, 20(3), 9-17. doi:10.3354/cr02009

Wibig, J. (1999). Precipitation in Europe in relation to circulation patterns at the 500 hPa level. *International Journal of Climatology*, 19(3), 253-269. doi:10.1002/(SICI)1097-0088(19990315)19:3<253::AID-JOC366>3.0.CO;2-O